



## Antioxidant, Antinociceptive and Anti-inflammatory Activities of Ethanolic Extract of Leaves of *Alocasia indica* (Schott.)

Mulla WA, Kuchekar SB<sup>1</sup>, Thorat VS, Chopade AR<sup>2</sup>, Kuchekar BS<sup>3</sup>

Government College of Pharmacy, Vidyanagar, Karad 415 124, <sup>1</sup>Allana College of Pharmacy, Pune-411 001, <sup>2</sup>Rajarambapu College of Pharmacy, Kasegaon, Taluka, Walwa District, Sangli- 415 404, Maharashtra, <sup>3</sup>MAEER's Maharashtra Institute of Pharmacy, Kothrud, Pune-411 038, India

*Address for correspondence:* Mr. Wahid A. Mulla; E-mail: wahidmulla@gmail.com

### ABSTRACT

Extracts obtained from the leaves of various *Alocasia* species have been used in India as folk remedy for the treatment of various inflammatory ailments including rheumatism and bruise. The ethanolic extract of leaves of *Alocasia indica* Schott. was evaluated by using different *in vitro* antioxidant models of screening like scavenging of 1, 1-diphenyl-2-picryl hydrazyl (DPPH) radical, nitric oxide radical, superoxide anion radical, and hydroxyl radical. The antinociceptive activity was tested by acetic acid-induced writhing response, hot plate method, and tail flick method in albino rats. The anti-inflammatory potential of gels of ethanolic extract has been determined by using carrageenan-induced paw edema assay, formalin-induced paw edema assay, arachidonic acid-induced ear edema assay, and xylene-induced ear edema assay. The extract showed remarkable antioxidant activity in all models, comparable to the standard reference drug ascorbic acid. The ethanolic extract of *Alocasia indica* and its gels produced dose-dependent antinociceptive and anti-inflammatory activity, respectively. This finding suggests that ethanolic extract of *A. indica* possess potent antinociceptive and anti-inflammatory activity possibly due to its free radical scavenging properties.

**Key words:** *Alocasia indica* Schott, antioxidant, antinociceptive, anti-inflammatory, ascorbic acid, diclofenac

**DOI:** 10.4103/0975-1483.63152

### INTRODUCTION

*Alocasia indica* Schott. is an indigenous herb belonging to family Araceae. Different parts of this plant are traditionally used in inflammation and in diseases of abdomen and spleen.<sup>[1]</sup> The juice of leaves of the plant is used as digestive, laxative, diuretic, astringent, and for the treatment of rheumatic arthritis.<sup>[2]</sup> It has hepatoprotective<sup>[3]</sup> and antifungal properties.<sup>[4]</sup> This plant contains flavonoids, cynogenetic glycosides, ascorbic acid, gallic acid, mallic acid, oxalic acid, alocaasin, amino acids, succinic acid, and  $\beta$ -lectines.<sup>[5]</sup> Since no scientific data are available to justify the traditional anti-inflammatory potential of the plant,

the present study was planned to validate the therapeutic use of this plant in treatment of inflammatory conditions.

### MATERIALS AND METHODS

#### Chemicals

All chemicals and solvents were of analytical grade and obtained from Loba Chemicals Ltd, Mumbai. 1, 1-diphenyl, 2-picryl hydrazyl (DPPH) was obtained from Sigma Chemicals, USA. The other chemicals used were sodium nitroprusside, sulphanilamide, *O*-phosphoric acid, naphthyl ethylene diamine dihydrochloride, trichloroacetic acid

(TCA), glacial acetic acid, nitroblue tetrazolium (NBT), ethylene diamine tetra acetic acid (EDTA), riboflavin, and Fe-EDTA. Carrageenan was obtained from Sigma-Aldrich, USA. Volini gel (Diclofenac sodium) was a gift sample from Ranbaxy Laboratories Ltd, Punjab. UV-Visible spectrophotometer (Shimadzu 1700) was used for recording the spectra.

### Experimental animals

Wistar albino rats weighing 175-225 g of either sex were obtained from Krishna Institute of Medical Sciences, Karad, Dist- Satara (Maharashtra) and were acclimatized for 10 days under standard housing conditions ( $24^{\circ} \pm 1^{\circ}\text{C}$ ; 45-55% RH with 12:12 h light/dark cycle). The animals had free access to rat food (Lipton Gold Mohr, India) and water. The animals were habituated to laboratory conditions for 48 h prior to the experimental protocol to minimize any nonspecific stress. The experimental protocol was approved by the Institutional Animal Ethics Committee by Government College of Pharmacy, Karad, India, and animals were maintained under standard conditions in the animal house approved by Committee for the Purpose of Control and Supervision on Experiments on Animals (CPCSEA).

### Plant material

Fresh leaves of *A. indica* Schott. (Araceae) collected from different places at Karad were authenticated at Botany Dept., Yashwantrao Chavan College of Science, Karad, and a voucher specimen was deposited at the Herbarium of institute.

### Preparation of extract

The ethanolic extract of leaves of *A. indica* was prepared by soxhletion. The powdered plant material (750 g) was repeatedly extracted in a round bottom flask with 2250 ml ethanol. The reflux time was 40 cycles for complete extraction. The extract was cooled at room temperature, filtered, and evaporated to dryness under reduced pressure in a rotary evaporator. The percentage yield was found to be 2.79 g. The ethanolic extract of *A. indica* was referred as AI.

### Phytochemical screening

Chemical tests were carried out on ethanolic extract of the plant for qualitative determination of phytochemical constituents.<sup>[6]</sup>

### In vitro antioxidant activity

*DPPH assay:* To 1 ml extract of different concentrations,

1 ml solution of DPPH (0.1 mM) was added. An equal amount of methanol and DPPH solution served as control. After 20 min of incubation in the dark, absorbance was measured at 517 nm.<sup>[7,8]</sup> Ascorbic acid was used as standard. The experiment was performed in triplicate and the percentage scavenging was calculated.

*Scavenging of nitric oxide radical:* Nitric oxide was generated from sodium nitroprusside and measured by Griss reaction.<sup>[9,10]</sup> Sodium nitroprusside (5 mM) in standard phosphate buffer saline solution (0.025 M, pH 7.4) was incubated with different concentrations of (200-1000  $\mu\text{g}/\text{ml}$ ) of the ethanolic extract dissolved in phosphate buffer saline (0.025 M, pH 7.4) and the tubes were incubated at  $25^{\circ}\text{C}$  for 5 h. Control experiments without test compounds but with equivalent amount of buffer were conducted in identical manner. After 5 h, 0.5 ml of solution was removed and diluted with 0.5 ml of Griss reagent (2 g of 1% sulphanilamide, 5 ml of 2% O-phosphoric acid, and 2 g of 0.1% naphthyl ethylene diamine dihydrochloride). The absorbance of chromophore formed during diazotization of nitrite with sulphanilamide and its subsequent coupling with naphthyl ethylene diamine was read at 546 nm. Ascorbic acid was used as standard. The experiment was performed in triplicate.

*Hydroxyl radical scavenging activity:* Extracts of different concentrations (200-1000  $\mu\text{g}/\text{ml}$ ) were taken in different test tubes and evaporated on water bath. To these, 1 ml of Fe-EDTA, 0.5 ml of EDTA, and 1 ml DMSO were added and the reaction was initiated by adding 0.5 ml ascorbic acid to each of the test tubes. Test tubes were capped tightly and heated in water bath at  $80-90^{\circ}\text{C}$  for 15 m. Then the reaction was terminated by addition of 1 ml of ice-cold TCA (17.5%, w/v) to all the test tubes and kept aside for 5 min. The formaldehyde formed was determined by adding 3 ml Nash reagent (75 g ammonium acetate, 3ml glacial acetic acid, 2 ml acetyl acetone was mixed and raised to 1 l with distilled water). This reaction mixture was kept aside for 15 min for color development.<sup>[11]</sup> Intensity of yellow color formed was measured spectrophotometrically at 412 nm against reagent blank. Ascorbic acid was used as standard. Percentage scavenging was calculated by comparison of the result of the samples, standard, with that of the blank.<sup>[12]</sup>

*Scavenging of superoxide radical by riboflavin--NBT system:* The assay was based on the capacity of the sample to inhibit blue formazon formation by scavenging the superoxide radicals generated in the riboflavin--NBT system. The reaction mixture contains 50 mM phosphate buffer pH 7.6, 20 g riboflavin, 12 mM NBT. Reaction was started by illuminating the test samples of the extract (200-1000  $\mu\text{g}/$

ml). The absorbance was measured at 590 nm. Ascorbic acid was used as positive control.<sup>[13,14]</sup>

The percentage scavenging of each extract in different concentrations for different *in vitro* models are calculated by following formula:

$$\% \text{ Scavenging} = \frac{\text{Absorbance of standard} - \text{absorbance of extract} \times 100}{\text{Absorbance of standard}}$$

IC<sub>50</sub> was calculated by using formulae:

$$b = \Sigma x.y / \Sigma x^2$$

$$a = \bar{y} - b \bar{x}$$

$$IC_{50} = a + b (50)$$

Where, *b* = regression coefficient of *x* on *y*; *a* = intercept of the line; *x* = concentration; and *y* = % scavenging;  $\bar{x}$  = mean of the concentration;  $\bar{y}$  = mean of % scavenging.

### Antinociceptive activity

**Writhing test:** Writhing was induced in rat (n = 6) by intraperitoneal injection (10 ml/kg) of 0.6% acetic acid. The number of writhings was counted over a 20 min period as previously reported.<sup>[15,16]</sup> Animals were treated through oral route 30 min before injection of acetic acid with ethanolic extract of AI (200 and 400 mg/kg) or acetylsalicylic acid (200 mg/kg). The control group received only vehicle (3 ml/kg) of 1% suspension of Tween-80.

**Tail immersion test:** Six rats were administered orally with vehicle (3 ml/kg), pentazocine (30 mg/kg), and ethanolic extract of AI (200 and 400 mg/kg). The distal part of the tails of the animals was immersed in hot water maintained at 55.0 ± 1.0 °C. The time taken to withdraw the tail was noted as reaction time.<sup>[17]</sup> A cut-off time of 10 s was maintained at 55 °C to prevent tissue damage. The reaction time was measured at 0, 15, 30, 45, and 60 min after treatment, respectively.

**Hot plate test:** Rats were placed on an aluminum hot plate kept at 55 ± 0.5 °C for a maximum time of 30 s.<sup>[18]</sup> Reaction time was recorded when the animals licked their fore- and hind paws and jumped; at before (0) and 15, 30, 45, and 60 min after intraperitoneal administration of 200 and 400 mg/kg of ethanolic extract of AI to different groups of six animals each. Morphine 10 mg/kg was used as the reference drugs.

### Anti-inflammatory activity

**Preparation of gels:** The gels of AI were prepared with carbopol-940 and PEG by cold method.<sup>[19]</sup> Different gels of AI (5%, 10%, and 20%) were formulated [Table 1]. The weighed amount of carbopol was placed in a beaker, and sufficient amount of water was added to it and kept in oven at 100 °C for 20 min to obtain a homogenous viscous mixture and cooled to room temperature with continuous stirring. Triethanolamine was added drop-wise with constant stirring. After formation of gel, weighed amount of AI was mixed in it.

**Carrageenan-induced paw edema assay:** The animals (n = 6) were divided into five groups. 0.5 g of AI gels (5%, 10% and 20%) and marketed formulation of diclofenac sodium (2%) were applied to the planter surface of the left hind paw by gently rubbing 50 times with the index finger.<sup>[20]</sup> The gel base without drug was applied in control group of rats by the same mode of application. After 2 h, the dose 0.1 ml of 1% carrageenan suspension (in sterile normal saline) was injected subplantarily into treated and control groups of rats. The paw volume was measured plethysmometrically (Ugo Basile, Italy) at 0 and 3 h after the carrageenan injection. The difference between the two readings was taken as the volume of edema, and the percentage anti-inflammatory activity was calculated using following equation:

$$\text{Percentage anti-inflammatory activity} = \frac{(V - V_i / V_i) \times 100}{V_i}$$

where *V* is the paw volume 3 h after the carrageenan injection and *V<sub>i</sub>* is the initial paw volume.

**Formalin-induced paw edema assay:** The same procedure as mentioned above in carrageenan-induced paw edema assay was followed except that acute inflammation was produced by administration of 20 µl formalin (2.5% in distilled water) into the subplanter area of right hind paw of rat.<sup>[21]</sup>

**Table 1: Formulation of gels of ethanolic extract of *Alocasia indica***

Ingredients (g)	Formulations (15 g)		
	5%	10%	20%
Ethanolic extract of <i>Alocasia indica</i>	0.75	1.50	3.00
carbopol-940	0.15	0.15	0.15
*PEG-4000	0.75	0.75	0.75
PEG-200	0.75	0.75	0.75
Ethanol	3.00	3.00	3.00
Triethanolamine	0.40	0.40	0.40
Methyl paraben	0.05	0.05	0.05
Propyl paraben	0.02	0.02	0.02
Distilled water	9.20	8.40	7.70

\*PEG: Polyethylene glycol.

**Arachidonic acid-induced ear edema assay:** The animals (n = 6) were divided into five groups. Samples were given 1 h prior to application of arachidonic acid.<sup>[22]</sup> One hour after topical application of AI gels (0.5 g) and marketed formulation of diclofenac sodium (2%), 2% arachidonic acid dissolved in acetone (0.02 ml/ear) was applied to anterior and posterior surfaces of right ear of rat. The gel base without drug was applied in the control group of rats by the same mode of application. The ear thicknesses were measured using a dial thickness gauge before and at 1 and 3 h after arachidonic acid treatment, and the differences in the thickness were calculated. The degree of ear swelling was expressed as an increase in ear thickness in mm.

**Xylene-induced ear edema assay:** The same procedure as mentioned above in arachidonic acid-induced ear edema assay was followed except that acute inflammation was produced by applying xylene to anterior and posterior surfaces of right ear of rat.<sup>[23]</sup>

### Statistical analysis

The statistical significance was assessed using one way of variance (ANOVA) followed by Bonferrini's multiple comparison tests. The values are expressed as mean  $\pm$  SE and  $P < 0.05$  was considered significant.

## RESULTS

### Preliminary phytochemical investigation

The preliminary phytochemical investigation of the ethanolic extract of *A. indica* showed that it contains flavonoids, cynogenetic glycosides, citric acid, ascorbic acid, and polyphenolic compounds.

### Antioxidant activity

Five concentrations ranging from 200 to 1000  $\mu\text{g/ml}$  of the ethanolic extract of *A. indica* were tested for their antioxidant potential using different *in vitro* models. It

was observed that free radicals were scavenged by test compounds at different concentrations. The maximum inhibitory concentration ( $\text{IC}_{50}$ ) in all models viz., DPPH, nitric oxide radical, superoxide, hydroxyl radical scavenging activity, were found to be 7.30, 10.97, 9.8, and 7.86  $\mu\text{g/ml}$ , respectively. The antioxidant model and percentage scavenging of each concentration of extract and standard are shown in Table 2.

### Antinociceptive activity

Oral administration of the ethanolic extract of AI (200 and 400 mg/kg) significantly ( $P < 0.05$ ) reduced the number of writhings induced by acetic acid in rat [Table 3]. The activity was comparable to that of acetylsalicylic acid (200 mg/kg, p.o.) used as a reference drug. Moreover, the extract induced protection in rat in tail immersion test [Table 4] that is comparable with the standard drug pentazocine (30 mg/kg, p.o.). The results of hot plate test presented in Table 5 showed that the i.p. administration of ethanolic extract of AI at the doses of 200–400 mg/kg significantly raised the pain threshold at observation time of 45 min in comparison with control ( $P < 0.001$ ).

### Stability of gels

Formulations exhibited desirable property and physical stability. AI was chemically stable in these formulations during 12-month storage at 4, 25, and 48 °C.

### Anti-inflammatory activity

Gels containing AI and marketed formulation of diclofenac sodium produced dose-dependent inhibition of carrageenan and formalin-induced paw edema as compared to the control ( $P < 0.05$ ) [Table 6]. In xylene-induced mouse ear edema assay gels containing AI and marketed formulation of diclofenac sodium significantly inhibited arachidonic acid and xylene-induced ear edema in dose dependent as compared to the control ( $P < 0.05$ ) [Table 7].

**Table 2: Antioxidant potential of ethanolic extract of *A. indica***

Different antioxidant models	% Scavenging at different concentration ( $\mu\text{g/ml}$ ) [values are mean of three replicates]							
	AA* ( $\mu\text{g/ml}$ )				Test extract: <i>A. indica</i> .			
	200	50	100	200	400	800	1000	IC <sub>50</sub>
DPPH**	88.75	40.24	44.62	53.25	61.39	63.46	73.48	7.30
Nitric oxide	74.54	34.74	53.44	63.48	71.59	73.51	84.09	10.97
Hydroxyl radical	62.35	43.24	55.37	66.15	57.23	59.28	67.96	9.8
Superoxide radical	88.25	47.26	52.18	65.71	68.12	52.84	77.17	7.86

\*Ascorbic acid; \*\*1,1-diphenyl, 2-picryl hydrazyl.

## DISCUSSION

Recent studies suggest that the inflammatory tissue damages are due to the liberation of reactive oxygen species from phagocytes invading the inflammation sites.<sup>[24-26]</sup> In addition to this, nitric oxide is also implicated in inflammation, cancer, and other pathological conditions.<sup>[27]</sup> Interactions between superoxide and nitric oxide regulate the vascular tone or inflammation.<sup>[28]</sup> The significant *in vitro* antioxidant activity of AI was in a concentration-dependent manner.

**Table 3: Effect of the ethanolic extract of leaves of *Alocasia indica* (AI) on acetic acid-induced writhing in rat**

Material	Dose (mg/kg)	No. of writhings	Inhibition (%)
Control	--	45.7 ± 4.2	
Acetylsalicylic acid	200	11.8 ± 2.4*	74.17
AI	200	21.5 ± 1.9**	52.95
	400	14.5 ± 1.8*	68.27

\* $P < 0.05$ , \*\* $P < 0.01$  vs. control, [Values are mean ± SE from six animals in each group.]

The ethanolic extract of AI at the doses of 200–400 mg/kg protected rat against both chemical- and thermal-induced noxious stimuli, which were evidenced from the acetic acid-induced writhing, tail immersion, and hot plate tests. Acetic acid induces writhing syndromes<sup>[15,16]</sup> and causes analgesia by releasing of endogenous substances, which then excite the pain nerve endings; the abdominal constriction is related to the sensitization of nociceptive receptors to prostaglandins.<sup>[29]</sup> Hot plate test was also assayed to characterize the analgesic activity of extract. It is possible that ethanol extract of AI extract exerts an analgesic effect probably by inhibiting the synthesis of prostaglandins. Variation in order of activity for ethanol extract of AI fractions in acetic acid-induced writhing, tail immersion, and hot plate tests indicated that the different constituents present in different fractions may be responsible for central and peripheral analgesia.

Carrageenan-induced edema is a biphasic response in which the involvement of the cyclo-oxygenase products of arachidonic acid metabolism and the production of reactive oxygen species are well established.<sup>[30]</sup> The first

**Table 4: Effect of the ethanolic extract of leaves of *Alocasia indica* (AI) on tail immersion method in rat**

Treatment	Dose (mg/kg)	Average tail withdrawing time (s)				
		0 min	15 min	30 min	45 min	60 min
Control	--	4.20 ± 0.11	4.65 ± 0.18	4.77 ± 0.17	4.77 ± 0.15	4.80 ± 0.18
Pentazocine	30	4.16 ± 0.20*	5.39 ± 0.21*	7.37 ± 0.45*	7.53 ± 0.29*	8.18 ± 0.27*
AI	200	4.25 ± 0.21*	4.53 ± 0.12**	5.27 ± 0.13*	5.35 ± 0.18**	5.27 ± 0.11**
	400	4.62 ± 0.16*	4.17 ± 0.28*	6.92 ± 0.48**	7.05 ± 0.28*	8.48 ± 0.27**

\* $P < 0.05$ , \*\* $P < 0.01$  vs. control, [Values are mean ± SE from six animals in each group.]

**Table 5: Effect of the ethanolic extract of leaves of *A. indica* (AI) on rat submitted to the hot plate test**

Treatment	Dose (mg/kg)	Time 0 (min)	Time 15 (min)	Time 30 (min)	Time 45 (min)	Time 60 (min)
Control	--	6.25 ± 0.35	6.32 ± 0.24	6.38 ± 0.31	7.15 ± 0.38	6.16 ± 0.54
Morphine	10	6.41 ± 0.35*	12.65 ± 0.48*	14.20 ± 0.36**	14.99 ± 0.81**	14.45 ± 0.56**
AI	200	5.76 ± 0.24**	10.16 ± 0.50*	13.21 ± 0.41*	14.17 ± 0.25**	12.16 ± 0.50*
	400	6.40 ± 0.48**	12.10 ± 0.31**	13.56 ± 0.73*	14.86 ± 0.72**	12.67 ± 0.78**

\* $P < 0.05$ , \*\* $P < 0.01$  vs. control, [Values are mean ± SE from six animals in each group.]

**Table 6: Effects of the *Alocasia indica* gels (AI: 5%, 10%, 20%) and marketed formulation of diclofenac sodium (DS: 2%) on carrageenan-induced rat paw edema and formalin-induced paw edema**

Group	Carrageenan-induced rat paw edema		Formalin-induced rat paw edema	
	Increase in paw volume (mean ± SEM) in ml	% Inhibition of paw edema	Increase in paw volume (mean ± SEM) in ml	% inhibition of paw edema
Control	0.67 ± 0.13	-	0.58 ± 0.12	-
AI (5%)	0.37 ± 0.07*	59.5	0.36 ± 0.05*	52.3
AI (10%)	0.32 ± 0.05*	66.6	0.31 ± 0.06*	58.1
AI (20%)	0.26 ± 0.03*	72.2	0.25 ± 0.02*	62.4
DS (2%)	0.20 ± 0.02*	83.1	0.19 ± 0.01**	73.7

\* $P < 0.05$ , \*\* $P < 0.01$  vs. control. [Values are mean ± SE from six animals in each group.]

**Table 7: Effects of the *Alocasia indica* gels (AI: 5%, 10%, 20%) and marketed formulation of diclofenac sodium (DS: 2%) on arachidonic acid-induced ear edema and xylene-induced ear edema**

Group	Arachidonic acid-induced ear edema		Xylene-induced ear edema	
	Increase in paw volume (mean $\pm$ SEM) in ml	% Inhibition of paw edema	Increase in paw volume (mean $\pm$ SEM) in ml	% Inhibition of paw edema
Control	0.27 $\pm$ 0.08	-	0.24 $\pm$ 0.05	-
AI (5%)	0.19 $\pm$ 0.06*	42.1	0.17 $\pm$ 0.06*	41.2
AI (10%)	0.15 $\pm$ 0.02*	45.3	0.14 $\pm$ 0.03*	44.4
AI (20%)	0.13 $\pm$ 0.03*	52.2	0.12 $\pm$ 0.03*	53.6
DS (2%)	0.09 $\pm$ 0.01**	63.5	0.08 $\pm$ 0.01**	61.1

\* $P < 0.05$ , \*\* $P < 0.01$  vs. control., [Values are mean  $\pm$  SE from six animals in each group.]

phase is mediated through the release of histamine, serotonin, and kinins, whereas the second phase is related to the release of prostaglandin oxygen-derived free radicals and production of inducible cyclo-oxygenase which peak at 3 h.<sup>[31]</sup> The AI extract produced dose-dependent and significant inhibition of carrageenan-induced paw edema comparable in magnitude with the inhibitory action of diclofenac. The formalin-induced paw edema assay defines distinctive biphasic nociceptive response termed neurogenic and inflammatory phases.<sup>[32]</sup> The ability of AI to have effect on both phases shows that it contains active anti-inflammatory principle acting both centrally and peripherally.

Another model of inflammation used is the ear edema induced by arachidonic acid and xylene, which serves to identify the mediators of inflammation of the lipoxygenase pathway.<sup>[33,34]</sup> Topical application of the gels of AI extract at the concentrations of 5%, 10%, and 20% inhibited the ear edema induced by arachidonic acid and xylene in dose-dependent manner similar to diclofenac. The results show that the AI acts on both the cyclo-oxygenase and lipoxygenase pathways.

Preliminary phytochemical screening showed that ethanolic extract of the leaves of *A. indica* contains flavonoid compounds. Flavonoids have been shown to possess various biological properties related to antioxidant, antinociceptive, and anti-inflammatory mechanisms by targeting reactive oxygen species and prostaglandins which are involved in the late phase of acute inflammation and pain perception.<sup>[35-37]</sup> It can be concluded that ethanolic extract of *A. indica* possess antioxidant, antinociceptive, and anti-inflammatory activities may be due to the presence of flavonoids and other polyphenolic moieties present in it, which seems to support the use of this plant in traditional medicine. Finally, the precise mechanism(s) and site(s) of this activity and the active constituent(s) involved are still to be determined in addition to toxicological studies.

## ACKNOWLEDGEMENT

The authors express their deep sense of gratitude to Dr S. B. Bhise, Principal Government College of Pharmacy, Karad, for providing the facilities to our research work. They also thank Professor S. K. Patil, Botany Department, Yashwantrao Chavan College of Science, Karad, Dist. - Satara (MH), for authentication work.

## REFERENCES

- Kirtikar KR, Basu BD. Ind Med Plants, 2<sup>nd</sup> ed. New Cannaught place, Dehradun: M/s Bishen Singh Mahendrapal Singh; 1975. p. 2617-8.
- Nadkarni KM. Indian Materia Medica. Mumbai: Popular Prakashan; 1996. p. 72-3.
- Mulla WA, Salunkhe VR, Bhise SB. Hepatoprotective activity of hydroalcoholic extract of leaves of *Alocasia indica*. Indian J Exp Biol 2009;47:316-21.
- Bhatt SK, Saxena VR. Antifungal activity of seeds of extract of *Alocasia indica*. Indian Drugs 1980;17:210.
- Prajapati ND. A Handbook of Medicinal plants. India: Agrobois Publication; 2003. p. 32.
- Khandelwal KR. Practical Pharmacognosy Techniques and Experiments. 9<sup>th</sup> ed. Pune: Nirali Prakashan; 2000.
- Sheejayan N, Rao MN. Nitric oxide scavenging by curcuminoids. J Pharm Pharmacol 1997;49:105-7.
- Shirwaikar A, Somashekar AP. Anti-inflammatory activity and free radical scavenging studies of *Aristolochia bracteolata* Lam. Indian J Pharm Sci 2003;65:68.
- Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JS, Tannenbaum SR. Analysis of nitrite and 15N in biological fluids. Anal Biochem 1982;126:131-8.
- Marcocci L, Maguire JJ, Droy- Lefaix MT and Packer L. The nitric oxide scavenging properties of Ginko biloba extract EGB 761. Biochem Biophys Res Commun 1994;201:748-55.
- Bhavani B, Pogozelski WK, Tullius TD. DNA strand breaking by hydroxyl radical is governed by the accessible surface area of the hydrogen atom of the DNA backbone. Pro Natl Acad Sci 1998;95:9738-43.
- Klein SM, Cohen G, Cedebaum AI. Production of formaldehyde during metabolism of dimethyl sulfoxide by hydroxyl radical generating system. Biochemistry 1991;20:6006-12.
- Robak J, Gryglewski RJ. Flavonoids are scavengers of superoxide anion. Biochem Pharmacol 1998;37:837.
- Beauchamp C, Fridovich I. Superoxide dismutase: Improved assays and an assay applicable to acryl amide gels. Anal Biochem 1971;44:276-87.
- Koster R, Anderson M, De Beer EJ. Acetic acid for analgesic screening. Fed Proc 1959;18:412.

## Antioxidant, Antinociceptive and Anti-Inflammatory Activities of *Alocasia indica* (Schott.)

16. Besra SE, Sharma RM, Gomes A. Anti-inflammatory effect of petroleum ether extract of leaves of *Litchi chinensis* Gaertn (Sapindaceae). *J Ethnopharmacol* 1996;54:1-6.
17. Vogel H, Vogel WH. Analgesic, anti-inflammatory and antipyretic activity in drug discovery and evaluation, pharmacological assays. New York: Springer; 1997. P. 360–418.
18. Vaz ZR, Mata LV, Calixto JB. Analgesic effect of the herbal medicine catuama in thermal and chemical models of nociception in mice. *Phytother Res* 1997;11:101–6.
19. Schmolka IS. Preparation and properties of pluronic F-127 gels for treatment for burns. *J Biomed Mater Res* 1972;6:571-82.
20. Winter CA, Risley EA, Nuss GW. Carrageenin-induced edema in hind paw of the rat as assay for anti-inflammatory drugs. *Proc Soc Exp Biol Med* 1962;11:544–7.
21. Chau TT. Analgesic testing in animal models, pharmacological methods in the control of inflammation. New York: Alan R Liss Inc.; 1989. p. 195.
22. Kim HK, Namgoong SY, Kim HP. Antiinflammatory activity of flavonoids: mice ear edema inhibition. *Arch Pharmacol Res* 1992;15:18–24.
23. Hosseinzadeh H, Haddadkhodaparast MH, Arash AR. Antinociceptive, anti-inflammatory and acute toxicity effects of *Salvia lerifolia* Benth seed extract in mice and rats. *Phytother Res* 2003;17:422–5.
24. Cross CE, Halliwell B, Borish ET, Pryor WA, Ames BN, Saul RL, et al. Oxygen radicals and human diseases. *Ann Intern Med* 1987;107:526-45.
25. Winrow VR, Winyard PG, Morris CJ, Blake DR. Free radicals in inflammation: second messengers and mediators of tissue destruction. *Br Med Bull* 1993;49:506-22.
26. Parke DV, Sapota A. Chemical toxicity and reactive oxygen species. *Int J Occup Med Environ Health* 1996;9:331–40.
27. Hemnani T, Parihar MS. Reactive oxygen species and oxidative DNA damage. *Indian J Physiol Pharmacol* 1998;42:440-52.
28. Conner EM, Grisham MB. Inflammation, free radicals and antioxidants. *Nutrition* 1996;12:274-7.
29. Vinegar R, Schreiber W, Hugo R. Biphasic development of carrageenin oedema in rats. *J Pharmacol Exp Ther* 1969;166:96-103.
30. Chen Q. Methodology in pharmacological study on Chinese materia medica. Beijing: 7 People's Medical Publishing House; 1993. p. 360.
31. Panthong A, Kanjanapothi D, Taesotikul T, Phankummoon A, Panthong K, Reutrakul V. Anti-inflammatory activity of methanolic extracts from *Ventilago harmandiana* Pierre. *J Ethnopharmacol* 2004;91:237–42.
32. Porro CA, Cavazzuti M. Spatial and temporal aspects of spinal cord and brainstem activation in the formalin pain model. *Prog Neurobiol* 1993;41:565–607.
33. Young JM, Young LM. In: Spector J, Back N, editors. Pharmacological methods in the control of inflammation. New York: Alan R. Liss; 1989. p. 215.
34. Matsuda R, Tanihata S. Suppressive effect of sialic acid on the prostaglandin E<sub>2</sub>-mediated oedema in carrageenin-induced inflammation of rat hind paws. *Nippon Yakurigaku Zasshi* 1992;99:363–72.
35. Hesham R, El-Seedi, Shgeru N. Chemistry of bioflavonoids. *Indian J Pharm Educ* 2007;39:172.
36. Di Carlo G, Mascolo N, Izzo AA, Capasso F. Flavonoids: old and new aspects of a class of natural therapeutic drugs. *Life Sci* 1999;65:337–53.
37. Rajnarayana K, Reddy MS, Chaluvadi MR, Krishna DR. Bioflavonoids classification, pharmacological, biochemical effects and therapeutic potential. *Indian J Pharmacol* 2001;33:2-16.

**Source of Support: Nil, Conflict of Interest: None declared.**